XALute

Watch out for these signs and symptoms!

As with any medicine, there may be side effects from your investigational study medicine. It's important you know what to watch out for and to report any symptoms or changes in how you feel to your study doctor right away. Prompt diagnosis and treatment can help side effects go away.



Feeling ill
- Fever
- Chills
- Headache
- Tiredness
- Feeling faint or dizzy

Eyes
- Swelling around eyes
- Yellowing of eyes

Heart
- Fast heart beat

Muscles and joints
- Aches or pains
- Weakness

Skin
- Rash
- Itchy or yellow skin
- Burning, prickling, or tingling

Legs and ankles
- Swelling

Neurological problems
- Loss of balance or coordination
- Shakiness of arms and/or legs
- Confusion or feeling disoriented
- Trouble speaking
- Memory loss
- Personality changes

Mouth and taste
- Change in taste, which may be foul, sour, salty, or metallic
- Blisters or sores
- Mouth pain

Throat
- Throat pain or swelling

Abdomen
- Nausea or vomiting
- Loss of appetite
- Abdominal pain

Genitals
- Swelling of scrotum

Report anything that is bothering you to your study doctor.
See the back of this page for more information.

XALURITAMIG

Cytokine release syndrome (CRS) is an important side effect to watch out for when taking xaluritamig. Symptoms of CRS include fever, chills, headache, low blood oxygen, changes in blood pressure, increased heart rate, nausea or vomiting, and increased liver enzymes.

Contact your study doctor or go to an emergency room if you have a fever (100.4°F/38.0°C or higher) or any other symptoms of CRS.

Localized inflammatory events such as muscle pain and inflammation are also associated with xaluritamig and may cause side effects that require prompt treatment to prevent worsening. Symptoms include:

- Muscle/joint pain and/or weakness
- Pain and/or swelling in the eyes, face, jaw, throat/mouth, or scrotum/genital area.

Other potential side effects of xaluritamig are described in greater detail in your Informed Consent Form.

CABAZITAXEL

The most common side effects are related to low blood cell counts (red blood cells, white blood cells, and platelets), diarrhea, tiredness, and lack of energy and strength.

ABIRATERONE ACETATE

The most common side effects include tiredness, joint pain, high blood pressure, nausea, swelling, hot flush, diarrhea, vomiting, upper respiratory tract infection (like a common cold), cough, and headache.

ENZALUTAMIDE

The most common side effects are muscle/joint pain, tiredness, hot flush, constipation, decreased appetite, diarrhea, high blood pressure, bleeding, fall, fracture, and headache.

These are not all the side effects. Your study doctor can give you a patient leaflet with more information about your investigational study medicine.

Scan the code or visit **www.xalutestudy.com** for additional study resources.

Pin Code: 0509